
How about a climate in 2070(doubling CO2)?


Rainfall Increase




Ranfall increases in blue region.


Average Temperature Increase is about 2.5

日本の真夏日日数と夏季平均気温の変化




日本の夏季豪雨日数と夏季平均降水量の変化






Changes in extreme temperatures


summer days Tmax>=25.0 [days/year]










-40-30-20-10-5-2-1 1 2 5 10 20 30 40



-40-	30-20-	10 -5	-2	-1	1	2	5	10	20	30	40








I 5W Ó SE 1ÓE 1∛E 2ĎE 2ĎE 3ĎE 3ĎE 4ĎE

-50-40-30-20-10-5-2 2 5 10 20 30 40 50

winter days Tmax<0.0 [days/year]




-50-40-30-20-10-5-2 2 5 1D 20 3D 40 50


Observation

Model(TL959)

(GPCP-1DD)

future change Daily precipitation Present






0.1 0.85


-0.5 -0.85 -0.1


Maximum number of consective dry days






An extremely intense typhoon appears in the future expt. (AS) with MIROC SST (max sfc wind 68.8 m/s, CP 877.6 hPa)







Tropical cyclone tracks




for Oct-Nov-Dec.

TC frequency as a function of peak wind intensity









Summary

- Recently, extreme events are increasing.
- Amount of loss is increasing!
- It is certain that probability of extreme events will increase in future!
- Development of a climate model will contribute to our adaptation for future climate change.